

On a differential algebraic approach of control observation problems

Sette Diop

► **To cite this version:**

| Sette Diop. On a differential algebraic approach of control observation problems. 2013. hal-00828592

HAL Id: hal-00828592

<https://hal-supelec.archives-ouvertes.fr/hal-00828592>

Submitted on 1 Jun 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On a Differential Algebraic Approach of Control Observation Problems

Sette Diop

Laboratoire des signaux & systèmes,
CNRS — Supélec — Université Paris Sud Orsay
Plateau de Moulon
91192 Gif sur Yvette cedex, France
diop@lss.supelec.fr

Abstract. Observation problems in control systems literature generally refer to problems of estimation of state variables (or identification of model parameters) from two sources of information: dynamic models of systems consisting in first order differential equations relating all system quantities, and online measurements of some of these quantities. For nonlinear systems the classical approach stems from the work of R. E. Kalman on the distinguishability of state space points given the knowledge of time histories of the output and input. In the differential algebraic approach observability is rather viewed as the ability to recover trajectories. This approach turns out to be a particularly suitable language to describe observability and related questions as structural properties of control systems. The present paper is an update on the latter approach initiated in the late eighties and early nineties. J. F. Pommaret, M. Fliess, S. T. Glad and the author.

Keywords: Control observation problems; State estimation; Differential algebraic decision methods; Differential algebraic geometry

1 Introduction

Observation problems in control systems literature generally refer to problems of estimation of state variables x from two sources of information: online measurements of external variables u and y , and first order dynamic models

$$\begin{cases} \dot{x} = f(t, u, x), \\ y = h(t, u, x), \end{cases} \quad (1)$$

relating x to u and y . See for instance [8, 10].

By using tools from differential algebraic geometry

- dynamic models are allowed to be implicit and of arbitrary order but restricted to be polynomial in variables and their derivatives,
- and may be considered more general situations of estimating one subset of system variables with respect to another subset of the system variables.

Specifically, given a dynamic system described by algebraic differential equations

$$\begin{cases} P_i(w, z, \zeta) = 0 & (i = 1, 2, \dots), \\ Q(w, z, \zeta) \neq 0, \end{cases} \quad (2)$$

one observation problem consists of the *online* estimation of $z(t) \in \mathbb{R}^\nu$ from the knowledge of the P_i 's and Q and time histories ($[t_0, t] \ni \tau \mapsto w(\tau) \in \mathbb{R}^\mu$) of w . Here the P_i 's and Q are differential polynomials in w, z and ζ , and ν, μ are natural integers. This problem is under investigation since the pioneering work of R. E. Kalman in the late fifties addressing its linear context. A complete non-linear answer is still lacking. A general approach consists of a two part theory: one of *observability*, that is, derivation of conditions on the P_i 's and Q guaranteeing the ability to some how estimate z from the supposedly known data, and the other part of the theory, the *observer design*, searches algorithms for such an estimation of z .

Though central the previous observation problem (observability and observer design) is not the only one. For instance, closely related to it, are two problems of *robustness* with respect to model and measurements uncertainties. Another observation problem with important practical application consists of determining subsets w of systems variables which make a given subset z observable.

Starting from the mid eighties (see [4, 5, 7, 15]) differential algebra and differential algebraic decision methods have been shown to provide a quite consistent language to describe some of these observation problems along with some of their solutions.

An account of this is proposed here. Some of the many open problems will be described.

2 The differential algebraic approach

A thorough introduction to the differential algebraic approach is available in [4]. For the sake of completeness the following definition is recalled from there.

A (*differential*) (*algebraic*) *system* \mathcal{X} with s variables, and with coefficients in a differential field \mathbf{k} is a *proper* differential quasi-affine variety $\mathcal{X} \subseteq \overline{\mathbf{k}}^s$ defined over \mathbf{k} where $\overline{\mathbf{k}}$ is a differential closure of \mathbf{k} . In observation problems, the system variable is partitioned into the *data*, or *observations*, $w = w_1, \dots, w_\mu$, the variable being observed (or estimated) $z = z_1, \dots, z_n$ and the remaining variables, ζ . In the classical observation problem, the data consist exclusively of (u, y) , the control u and the measurements y . When the variable ζ is present, the projection $\mathcal{X}_{w,z}$ of \mathcal{X} along the variable ζ is considered. It is the set of elements $(\overline{w}, \overline{z}) \in \overline{\mathbf{k}}^\mu \times \overline{\mathbf{k}}^n$ such that there is at least $\overline{\zeta}$ such that $(\overline{w}, \overline{z}, \overline{\zeta}) \in \mathcal{X}$.

In terms of equations, previously defined systems are those described by

$$\begin{cases} P_i(w, z, \zeta) = 0, & i = 1, 2, \dots, \\ Q(w, z, \zeta) \neq 0, \end{cases} \quad (3)$$

where the P_i 's and Q are finitely many polynomials in w, z, ζ and their derivatives.

For a system \mathcal{X} the variable z is said to be (*algebraically*) *observable with respect to w* if the projection map $\pi : \mathcal{X}_{w,z} \rightarrow \mathcal{X}_w$ (sending every trajectory (\bar{w}, \bar{z}) of $\mathcal{X}_{w,z}$ into the corresponding observation \bar{w}) is *generically finite*. If z is observable with respect to w then the degree of π is called the *observability degree* of z with respect to w , and is denoted by $d_w^\circ z$. The variable z is said to be *rationally* observable with respect to w if it is observable with respect to w with observability degree one. State systems of the form (1) are said to be *observable* if x is observable with respect to (u, y) .

It was first proved in [5] (see [4] for more details) that the previous definition has a differential algebraic translation, namely: z is observable with respect to w iff z is algebraic over $\mathbf{k}\langle w \rangle$, that is, for each component, z_i of z there is a polynomial equation

$$H_i(z_i, w, \dot{w}, \dots) = 0 \quad (4)$$

in z_i , and finitely many time derivatives of the data w , with coefficients in \mathbf{k} .

The reader is referred to [4] more details on differential algebraic geometry terms or notations used here without explanations.

3 How does it compare to the classical theory?

Formal definitions of observability can be found in [8, 10] for instance.

For linear state systems

$$\begin{cases} \dot{x} = Fx + Gu, \\ y = Hx + Eu, \end{cases} \quad (5)$$

the answer to the question is that algebraically observability of x with respect to (u, y) is *equivalent* to the classical Kalman definition of observability of system 5. The proof of this is as follows.

As is well known system 5 is observable in the classical sense iff $\text{rk}_{\mathbf{R}} \mathcal{O}(F, H) = n$, where n is the number of components of the state, x , and where

$$\mathcal{O}(F, H) = \begin{pmatrix} H \\ HF \\ \vdots \\ HF^{n-1} \end{pmatrix}.$$

Now the following equalities

$$\begin{aligned} Hx &= y - Eu && = z_0, \\ HFx &= \dot{z}_0 - HG u && = z_1, \\ HF^2x &= \dot{z}_1 - HFG u && = z_2, \\ &\vdots \\ HF^{n-1}x &= \dot{z}_{n-2} - HF^{n-2}Gu = z_{n-1}, \end{aligned} \quad (6)$$

result from the equations of system 5. The reader will notice that there are written such that only supposedly differentiable quantities are differentiated. They may be rewritten as

$$\mathcal{O}(F, H) x = \begin{pmatrix} z_0 \\ z_1 \\ z_2 \\ \vdots \\ z_{n-1} \end{pmatrix}. \quad (7)$$

Therefore if system 5 is observable in the classical sense then it is so in the algebraic sense. The converse follows from Corollary 1.

The equivalence between the algebraic and the classical definitions of observability for the class of systems 5 was first mentioned in § 5.1.2 of [16] but it was not shown how the Cayley-Hamilton theorem which was invoked does complete the proof.

It is worth emphasizing the fact that the algebraic definition of systems applies without any change to so-called *implicit* or *descriptor* linear systems

$$\begin{cases} M \dot{x} = F x + G u, \\ y = H x + E u, \end{cases}$$

where the matrix M is *singular*. Compare to [2] and references therein.

The largest class of systems where algebraic and classical observability may be compared is the class of rational state systems

$$\begin{cases} \dot{x}_i = \frac{p_i(u, x)}{q_i(u, x)} \quad (1 \leq i \leq n), \\ y_j = \frac{f_j(u, x)}{g_i(u, x)} \quad (1 \leq j \leq p), \end{cases} \quad (8)$$

where u stands for u_1, u_2, \dots, u_m , and p_i, q_i, f_i and g_i are differential polynomials of *order zero* in x with coefficients in $\mathbf{k} = \mathbb{R}$, the algebraic observability (of x with respect to (u, y)) is equivalent to the *generic local observability* of the system as defined in [10]. This was first obtained in [6].

4 Partial answers to some observation problems

Among all benefits of the differential algebraic approach to observation problems, application of decision methods is perhaps the most appealing.

4.1 Computing

One of the most used estimation algorithms, the Kalman filter (or its extended form), is often applied in practice without prior asserting the validity of its

conditions. The reason of this is that there is no systematic method for such a verification.

The differential algebraic theory of observability is constructive *in principle*.

Most of the decision methods which may be used are already described in [4]. This is the case for general polynomial systems

$$P_i(w, z, \zeta) = 0, \quad i = 1, 2, \dots \quad (9)$$

with *constant* coefficients. For rational state systems 8 the observability test is formally similar to the Jacobian rank condition which is found in [10].

For polynomial state systems with *nonconstant* coefficients let \mathbf{k} be an *ordinary* differential field (*not necessarily of constants*). Let \mathcal{X} be

$$\begin{cases} \dot{x}_i = f_i(u, x) & (1 \leq i \leq n), \\ y_j = h_j(u, x) & (1 \leq j \leq p), \end{cases} \quad (10)$$

where the f_i 's and h_j 's are (nondifferential) *polynomials* in their arguments with coefficients in \mathbf{k} . Let

$$\begin{aligned} P_i(U, X, Y) &= X_i^{(1)} - f_i(U, X) & (1 \leq i \leq n); \\ P_{n+j}(U, X, Y) &= h_j(U, X) - Y_j & (1 \leq j \leq p). \end{aligned}$$

be the differential polynomials defining \mathcal{X} . Let $\sigma : \mathbf{k}\{U, Y\} \rightarrow \mathbf{k}\{u, y\}$ be the substitution map which sends U into u and Y into y , where $\mathbf{k}\{u, y\}$ is the differential \mathbf{k} -subalgebra of $\mathbf{k}\{u, x, y\}$ generated over \mathbf{k} by u and y . Let P be in $\mathbf{k}\{U, X, Y\}$ and P^σ denote the element of $\mathbf{k}\{u, y\}\{X\}$ obtained by regarding P as a differential polynomial in X with coefficients in $\mathbf{k}\{U, Y\}$ and by applying σ to each of these coefficients, and let $\mathbf{I}(\mathcal{X})^\sigma$ stand for the differential ideal of $\mathbf{k}\{u, y\}\{X\}$ consisting of P^σ ($P \in \mathbf{I}(\mathcal{X})$). The ideal of definition, \mathfrak{a} , of $\mathbf{k}\langle u, y \rangle(x)$ over $\mathbf{k}\langle u, y \rangle$ is equal to $\mathbf{I}(\mathcal{X})^\sigma \cap \mathbf{k}\langle u, y \rangle[(X_i)_{1 \leq i \leq n}]$. Note that the set \mathcal{A} consisting of the P_i ($1 \leq i \leq n$) form an *autoreduced set* with respect to any *ranking* of $\mathbf{k}\{U, X, Y\}$ such that U, Y and their derivatives all are lower than X .

Let us now inductively define some polynomials which will turn out to be generators of the ideal $\mathbf{I}(\mathcal{X})^\sigma \cap \mathbf{k}\langle u, y \rangle[(X_i)_{1 \leq i \leq n}]$ of $\mathbf{k}\langle u, y \rangle[(X_i)_{1 \leq i \leq n}]$.

Starting with

$$Q_i(U, X, Y) = P_{n+i}(U, X, Y) \quad (1 \leq i \leq p),$$

then let Q_{p+i} be the *remainder* of the derivative of Q_i ($1 \leq j \leq n$) with respect to the previously mentioned autoreduced set, \mathcal{A} . The polynomial Q_{p+i} is merely the derivative of Q_i in which $X_j^{(1)}$ is eliminated by substituting $P_j + f_j$ for $X_j^{(1)}$ ($1 \leq j \leq n$) (The linear combination of P_j ($1 \leq j \leq n$) which appears reduces to zero when the remainder is taken, so that it can be ignored.) Explicitly, Q_{p+i} is as follows

$$\begin{aligned} Q_{p+i} &= \sum_{1 \leq j \leq n} \frac{\partial Q_i}{\partial U_j} U_j^{(1)} + \sum_{1 \leq j \leq n} \frac{\partial Q_i}{\partial X_j} f_j - Y_i^{(1)} \\ &+ Q_i \bullet \quad (1 \leq i \leq p), \end{aligned}$$

where the notations

$$P_{\bullet} \equiv P_{(1)}, \quad P_{(2)}, \quad \dots \tag{11}$$

for a differential polynomial P stand for the differential polynomials obtained by replacing the coefficients of P by their respective derivatives respectively at order 1, 2, etc.

Note that this formula is nothing but a counterpart of Lie derivatives: Authors usually consider the functions h_j as free of u and the functions f_i and h_j as with *constant* coefficients so that in the left hand side of the latter equation the first sum as well as the last term are absent.

This construction of Q_{p+i} from Q_i is iterated in order to get Q_{2p+i} ($1 \leq i \leq p$) as the remainder of the derivative of Q_{p+i} ($1 \leq i \leq p$). And so on.

By their definition,

$$Q_i^\sigma \in \mathbf{I}(\mathcal{X})^\sigma \cap \mathbf{k}\langle u, y \rangle \left[(X_j)_{1 \leq j \leq n} \right] \quad (i \in \mathbb{N}).$$

Conversely, let P

$$P \in \mathbf{I}(\mathcal{X})^\sigma \cap \mathbf{k}\langle u, y \rangle \left[(X_j)_{1 \leq i \leq n} \right].$$

As an element of $\mathbf{I}(\mathcal{X})$, P may easily be written in the form

$$P = \sum_{1 \leq i \leq n, j \in \mathbb{N}} A_{i,j} P_i^{(j)} + \sum_{i \in \mathbb{N}} B_i Q_i.$$

where B_i ($i \in \mathbb{N}$) are in

$$\mathbf{I}(\mathcal{X}) \cap \mathbf{k}\{U, Y\} \left[(X_i)_{1 \leq i \leq n} \right].$$

Since the differential ideal of $\mathbf{k}\{U, X, Y\}$ generated by P_i ($1 \leq i \leq n$) has no nonzero element in common with $\mathbf{I}(\mathcal{X}) \cap \mathbf{k}\{U, Y\} \left[(X_j)_{1 \leq i \leq n} \right]$ (this results from an obvious degree argument), the first sum in the previous equality must be zero.

This ends the proof that Q_i^σ ($i \in \mathbb{N}$) form a basis of \mathfrak{a} .

Lemma 1. *A set of generators of the ideal of definition of $\mathbf{k}\langle u, y \rangle(x)$ over $\mathbf{k}\langle u, y \rangle$ is given by*

$$\begin{aligned} & Q_1(u, X, y), Q_2(u, X, y), \dots, Q_p(u, X, y), \\ & Q_{p+1}(u, X, y), Q_{p+2}(u, X, y), \dots, Q_{2p}(u, X, y), \\ & \dots \end{aligned}$$

In addition, it comes from the Hilbert basis theorem that only finitely many Q_i suffice to generate the ideal \mathfrak{a} . That is, there is some μ in \mathbb{N} such that the first μ rows of the previous list of $Q_i(u, X, y)$ generate the ideal of definition of

$\mathbf{k}\langle u, y \rangle(x)$ over $\mathbf{k}\langle u, y \rangle$. According to Theorem 16 of [4], the observability of \mathcal{X} is equivalent to the fact that the $\mathbf{k}\langle u, y \rangle(x)$ -matrix

$$\left[\frac{\partial Q_i}{\partial X_j} (u, x, y) \right]_{\substack{1 \leq i \leq \mu p \\ 1 \leq j \leq n}}$$

is of rank n .

Now it is a basic fact that the above rank is equal to the rank of the first np rows.

Corollary 1. *If \mathcal{X} possesses a state description as above, then \mathcal{X} is observable if, and only if, the following $\mathbf{k}\langle u, y \rangle(x)$ -matrix*

$$\left[\frac{\partial Q_i}{\partial X_j} (u, y, x) \right]_{\substack{1 \leq i \leq np \\ 1 \leq j \leq n}}$$

(which is formally the counterpart of the matrix of Lie derivatives which appears in the Hermann-Krener observability Jacobian rank condition) is of rank n .

The main difference between this rank condition and the one in [10] is that the rank is not over \mathbf{k} (which is usually \mathbb{R}) but over a much bigger field (and, here the rank condition is a necessary and sufficient condition).

For arbitrary systems observability tests resort on decision methods such as characteristic set of the defining differential ideal of \mathcal{X} . See [4] for more details. Very promising, Thomas decomposition was also proposed as decision methods for the same tests, see [1, 14].

5 Regular observability

The notion of regular observability refers to the classical one of *universal inputs* as thoroughly treated in [19]. Bad inputs (as opposed to universal ones) are supposed to occlude the functioning of online estimation schemes when they happened to be applied to a system. The present differential algebraic approach has brought a new light to this notion of singularity of the observability property. Here is an abstract of the result which may be found in [4] in more details.

Let \mathcal{X} be a system with variables w , z , and ζ , and with coefficients in \mathbf{k} . It is a matter of fact that, when z is observable with respect to w then for special observations \bar{w} , $\pi^{-1}(\bar{w})$ may contain infinitely many elements, leading to a singularity of the generic notion of observability. Here π is the projection map of § 2. An example of such situations is the following

$$\begin{cases} \dot{x}_1 = x_1 x_2, \\ \dot{x}_2 = u + x_2, \\ y = x_1. \end{cases} \quad (12)$$

x is observable with respect to u, y since

$$x_1 = y \quad \text{and} \quad x_2 = \frac{\dot{y}}{y}.$$

But in practice, in any time interval where y is identically zero (or, merely, small), the observability of x_2 is singular in the sense that it is lost.

An observation $\bar{w} \in \mathcal{X}_w$ is said to be *singular* for the observation of z with respect to w if $\pi^{-1}(\bar{w})$ is *infinite*. Observations $\bar{w} \in \mathcal{X}_w$ which are not singular are called *regular*. The variable z is said to be *regularly observable* with respect to w if there is no singular observation for its observability with respect to w .

The best result obtained in this approach reads as follows.

Theorem 1. *Let \mathcal{X} be a system with variables w, z , and ζ , and with coefficients in \mathbf{k} . The variable z is regularly observable with respect to w if z is primitive over $\mathbf{k}\{w\}$.*

5.1 Sensor selection

Given a dynamic system with differential field extension

$$\mathbf{k}\langle u, z \rangle$$

with input u and latent variable z the sensor selection problem consists of the selection of sensors which endow the system with some properties. Among all such desirable properties is the basic one of observability. In this section partial answers to the following questions will be provided.

1. What is the minimal number of sensors that make the dynamics observable?
2. When the sensors are bound to measure state components, what is their minimum number?
3. How may the *observability margin* be improved by selecting the sensors?

Let y denote an arbitrary output of the system. By definition, an output, y , is componentwise algebraic over $\mathbf{k}\langle u, z \rangle$. An output makes the dynamics observable if

$$d_{\mathbf{k}\langle u, z, y \rangle}^{\circ} \mathbf{k}\langle u, y \rangle = 0,$$

that is, if each component of z is algebraic over $\mathbf{k}\langle u, y \rangle$. Clearly, $y = z$ is an output which makes the system observable. Let n denote the number of components of z , and

$$\mathbf{I} = \left\{ p \in \mathbb{N} : \exists y_1, y_2, \dots, y_p \in \mathbf{k}\langle u, z \rangle, d_{\mathbf{k}\langle u, z, y \rangle}^{\circ} \mathbf{k}\langle u, y \rangle = 0 \right\}$$

The set \mathbf{I} is the one of integers p such that there exists an output y with p components which makes the system observable.

It is a nonempty (since $n \in \mathbf{I}$) subset of \mathbb{N} . Therefore, \mathbf{I} contains a smallest element which is precisely the minimal number of sensors which make the system observable.

The sensor selection problem characterizing this minimum number, p , of sensors is an open problem.

Later in this section it is shown that the minimum number of sensors is 1 for rational state systems, providing a partial answer to Question 1 above.

Next, about Question 2, what if the output y is chosen as a subset of the components of z , instead of vector rational function of u and z ? A complete but trivial, inelegant, and computationally costly answer consists of performing the $2^n - 1$ observability tests!

Up to the knowledge of the author there is no partial contribution to Question 3.

Back to Question 1, here is the surprising answer for the class of rational state systems.

Theorem 2. *Let the state of a system \mathcal{X} be given by*

$$\dot{x} = f(u, x) \quad (13)$$

with a vector rational function f of input u , state x , and with coefficients in a differential field \mathbf{k} . Let m and n be the respective numbers of components of u and x . Let \mathbf{K} be a differential extension field of \mathbf{k} . If \mathbf{K} contains nonconstants then there always is a scalar output

$$y = \sum_{i=1}^n \alpha_i x_i \quad (14)$$

with $\alpha_1, \alpha_2, \dots, \alpha_n$ in \mathbf{K} , which makes x observable with respect to (u, y) . Moreover, for y as in (14) to make \mathcal{X} observable it is sufficient that the associated α 's be linearly independent over the subfield of constants of \mathbf{K} .

Proof. Let $y^{[n]}$ denote the vector

$$y^{[n]} = \begin{pmatrix} y \\ \dot{y} \\ \vdots \\ y^{(n-1)} \end{pmatrix}.$$

By Corollary 1, for the output (14) to make x observable with respect to (u, y) it is necessary, and sufficient, that the Jacobian matrix of $y^{[n]}$ with respect to x

$$\frac{\partial y^{[n]}}{\partial x'} = \begin{pmatrix} \frac{\partial y}{\partial x'} \\ \frac{\partial \dot{y}}{\partial x'} \\ \vdots \\ \frac{\partial y^{(n-1)}}{\partial x'} \end{pmatrix} = \begin{pmatrix} \frac{\partial y}{\partial x_1} & \frac{\partial y}{\partial x_2} & \cdots & \frac{\partial y}{\partial x_n} \\ \frac{\partial \dot{y}}{\partial x_1} & \frac{\partial \dot{y}}{\partial x_2} & \cdots & \frac{\partial \dot{y}}{\partial x_n} \\ \vdots & \vdots & \cdots & \vdots \\ \frac{\partial y^{(n-1)}}{\partial x_1} & \frac{\partial y^{(n-1)}}{\partial x_2} & \cdots & \frac{\partial y^{(n-1)}}{\partial x_n} \end{pmatrix}$$

be of rank n over $\mathbf{K}\langle u, y \rangle(x)$ where the complete system is considered as with coefficients in $\mathbf{K} \supseteq \mathbf{k}$.

Now y may be written as

$$y = \alpha' x$$

where

$$\alpha = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix}.$$

The first row of $\partial y^{[n]} / \partial x'$ is merely

$$\frac{\partial y}{\partial x'} = \alpha'.$$

The second row of the same matrix is

$$\frac{\partial \dot{y}}{\partial x'} = \dot{\alpha}' + \alpha' \frac{\partial f(u, x)}{\partial x'}.$$

Therefore,

$$\text{rk}_{\mathbf{k}\langle u, y \rangle(x)} \frac{\partial y^{[n]}}{\partial x'} = \text{rk}_{\mathbf{k}\langle u, y \rangle(x)} \begin{pmatrix} \alpha' \\ \dot{\alpha}' \\ \frac{\partial \dot{y}}{\partial x'} \\ \vdots \\ \frac{\partial y^{(n-1)}}{\partial x'} \end{pmatrix}$$

by substituting the linear combination

$$\frac{\partial \dot{y}}{\partial x'} - \frac{\partial y}{\partial x'} \frac{\partial f(u, x)}{\partial x'} = \dot{\alpha}'$$

over $\mathbf{k}\langle u, y \rangle(x)$ of the first two rows for the second row of $\partial y^{[n]} / \partial x'$.

More generally, by the Leibniz formula,

$$y^{(i)} = \alpha^{(i)'} x + \sum_{j=1}^i \binom{i}{j} \alpha^{(i-j)'} x^{(j)},$$

and by the fact that $x^{(j)}$ is in $\mathbf{k}\langle u \rangle(x)$ for all $j \geq 1$, it is clear that

$$\text{rk}_{\mathbf{k}\langle u, y \rangle(x)} \frac{\partial y^{[n]}}{\partial x'} = \text{rk}_{\mathbf{k}\langle u, y \rangle(x)} \begin{pmatrix} \alpha' \\ \dot{\alpha}' \\ \ddot{\alpha}' \\ \vdots \\ \alpha^{(n-1)'} \end{pmatrix}$$

by an immediate induction on the row's index of the Jacobian matrix.

Now note that the matrix

$$\mathcal{W}(\alpha) = \begin{pmatrix} \alpha' \\ \dot{\alpha}' \\ \vdots \\ \alpha^{(n-1)} \end{pmatrix}$$

is square, and of order n , and does not involve neither u nor x . Therefore it is of rank n over $\mathbf{K}(u, y)(x)$ if, and only, its determinant is nonzero.

Next note that the determinant of $\mathcal{W}(\alpha)$ is simply a differential polynomial in $\alpha_1, \alpha_2, \dots, \alpha_n$ with coefficients in the field of constants of \mathbf{K} . Then the following theorem is used.

Theorem 3. *If G is a nonzero differential polynomial in n indeterminates with coefficients in a differential field containing nonconstant elements then G possesses a zero (z_1, \dots, z_n) over \mathbf{K} .*

For a proof see [12, 13, 17, 18] for instance.

This terminates the proof of the first assertion in the theorem.

The second assertion follows from the following. The n elements $\alpha_1, \alpha_2, \dots, \alpha_n$ of \mathbf{K} are said to be linear dependent over constants if there is a nontrivial relation

$$c_1 \alpha_1 + c_2 \alpha_2 + \dots + c_n \alpha_n = 0$$

with constant coefficients. It is a classical result that

Theorem 4. *$\alpha_1, \alpha_2, \dots, \alpha_n$ are linearly dependent over constants if, and only if, the Wronskian matrix $\mathcal{W}(\alpha)$ is singular.*

For a proof see the same references [12, 13, 17, 18] for instance. This ends the proof of the theorem.

6 Some of the questions without partial answers

The following is no way an exhaustive list of open problems. It is simply believed that the reader may be inspired to contribute to their solution. One of the most challenging open question is actually of a foundation level.

6.1 A foundation problem

Observation problems are basically encountered in engineering practices where *real problems* often refer to *real valued* parameters and variables. But as the reader has already noticed the present differential algebraic geometry approach has recourse to so-called *differential closures* of found fields. Basically such fields are complex ones. For more details the reader may refer to section 4.1 of [4].

6.2 Robustness

Keeping in mind that engineering observation problems deal with systems which may be inaccurately known. The most favorable lack of information is actually that of true values of parameters or coefficients: in other words, the form and orders and dimensions of the equations are exactly known, only actual parameter values are uncertain.

The question is then how observability and other observation problems assertions behave in the presence of parameter uncertainties?

For linear systems 5 a notion of *observability margin* may be defined characterizing the *distance* of a given system to unobservable ones. This generally uses matrix tools such as singular values.

For general systems the question is clearly related to the so-called field of *decisions methods for approximate systems* as tackled in [3, 11] and many other papers.

6.3 Decision methods problems

Among decision methods capable of dealing with real example such as those one may find in biotechnology are also wanted. In this vein, there is a question with practical importance: Given an observable variable z with respect to w , what is the minimal order of derivatives of w involved in the observability of z with respect to w ?

References

- [1] T. BÄCHLER, V. GERDT, M. LANGE-HEGERMANN, and D. ROBERTZ, *Thomas decomposition of algebraic and differential systems*, Computer Algebra in Scientific Computing, ed. by V. P. GERDT, W. KOEPF, E. W. MAYR, and E. V. VOROZHTSOV, Lect. Notes Comput. Sci. vol. 6244, Springer, Berlin, Heidelberg, 2002, pp. 31–54 DOI: [10.1007/978-3-642-15274-0_4](https://doi.org/10.1007/978-3-642-15274-0_4).
- [2] F. J. BEJARANO, T. FLOQUET, W. PERRUQUETTI, and GANG ZHENG, *Observability and detectability analysis of singular linear systems with unknown inputs*, Proceedings of the IEEE Conference on Decision and Control, IEEE Press, New York, 2011, pp. 4005–4010 DOI: [10.1109/CDC.2011.6161325](https://doi.org/10.1109/CDC.2011.6161325).
- [3] G. CHÈZE and A. GALLIGO, From an approximate to an exact absolute polynomial factorization, *J. Symbolic Comput.* **41**(2006), 682–696, DOI: [10.1016/j.jsc.2005.11.004](https://doi.org/10.1016/j.jsc.2005.11.004).
- [4] S. DIOP, *From the geometry to the algebra of nonlinear observability*, Computer Algebra in Scientific Computing, ed. by V. P. GERDT, W. KOEPF, E. W. MAYR, and E. V. VOROZHTSOV, Lect. Notes Comput. Sci. vol. 6244, Springer, Berlin, Heidelberg, 2002, pp. 305–345 DOI: [10.1142/9789812778079_0012](https://doi.org/10.1142/9789812778079_0012).

- [5] S. DIOP and M. FLIESS, *On nonlinear observability*, Proceedings of the European Control Conference, ed. by C. COMMAULT, D. NORMAND-CYROT, J. M. DION, L. DUGARD, M. FLIESS, A. TITLI, G. COHEN, A. BENVENISTE, and I. D. LANDAU, Hermès, Paris, 1991, pp. 152–157.
- [6] S. DIOP and Y. WANG, *Equivalence between algebraic observability and generic local observability*, Proceedings of the IEEE Conference on Decision and Control, vol. 3, IEEE Press, New York, 1993, pp. 2864–2865 DOI: [10.1109/CDC.1993.325718](https://doi.org/10.1109/CDC.1993.325718).
- [7] M. FLIESS, *Quelques remarques sur les observateurs non linéaires*, Proceedings Colloque GRETSI Traitement du Signal et des Images, GRETSI, 1987, pp. 169–172.
- [8] J. P. GAUTHIER and I. A. K. KUPKA, *Deterministic Observation Theory and Applications*, Cambridge University Press, Cambridge, United Kingdom, 2001. DOI: [10.1017/CB09780511546648](https://doi.org/10.1017/CB09780511546648).
- [9] V. P. GERDT, W. KOEPF, E. W. MAYR, and E. V. VOROZHTSOV, eds. *Computer Algebra in Scientific Computing* vol. 6244 Lect. Notes Comput. Sci. Springer, Berlin, Heidelberg, 2010.
- [10] R. HERMANN and A. J. KRENER, Nonlinear controllability and observability, *IEEE Trans. Automat. Control*, **22**(1977), 728–740, DOI: [10.1109/TAC.1977.1101601](https://doi.org/10.1109/TAC.1977.1101601).
- [11] E. KALTOFEN, J. P. MAYE, Z. YANG, and L. ZHI, Approximate factorization of multivariate polynomials using singular value decomposition, *J. Symbolic Comput.* **43**(2008), 359–376, DOI: [10.1016/j.jsc.2007.11.005](https://doi.org/10.1016/j.jsc.2007.11.005).
- [12] I. KAPLANSKY, *An Introduction to Differential Algebra. Second Edition*, Hermann, Paris, 1976.
- [13] E. R. KOLCHIN, *Differential Algebra and Algebraic Groups*, Academic Press, New York, 1973.
- [14] M. LANGE-HEGERMANN and D. ROBERTZ, *Thomas decompositions of parametric nonlinear control systems*, Technical report, 2012 URL: <http://arxiv.org/abs/1212.0377>,
- [15] J. F. POMMARET, Géométrie différentielle algébrique et théorie du contrôle, *C. R. Acad. Sci. Paris Sér. I*, **302**(1986), 547–550.
- [16] J. F. POMMARET, *Partial Differential Control Theory. Volume II: Control Systems*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2001.
- [17] J. F. RITT, *Differential Algebra*, American Mathematical Society, Providence, 1950.
- [18] A. SEIDENBERG, Some basic theorems in differential algebra (characteristic p , arbitrary), *Trans. Amer. Math. Soc.* **73**(1952), 174–190, DOI: [10.1090/S0002-9947-1952-0049174-X](https://doi.org/10.1090/S0002-9947-1952-0049174-X).
- [19] H. J. SUSSMANN, Single-input observability of continuous-time systems, *Math. Systems Theory*, **12**(1979), 371–393, DOI: [10.1007/BF01776584](https://doi.org/10.1007/BF01776584).